



# CLEANASCITE™ & BINDPRO™

## LIPID/PROTEIN REMOVAL & CELL RESPONSE APPLICATIONS

FEBRUARY 24, 2020

BIOTECH SUPPORT GROUP  
Sample Prep that Matters

### Introduction


The “omics” revolution demanded new and different sample prep separations that were not efficiently performed by conventional technologies. While effective for many applications, these tools were not efficient for “omics” sample preparation, as throughput, economy and simplicity are especially required. Furthermore, these same separation tools often denatured proteins which limited their use in applications which demanded the measurement of function, structure or bio-activity.

For these reasons, BSG has been dedicated to create new methods and applications to drive efficient workflows and better data quality for all proteomic and biomarker analyses. Of special importance is the value created when certain families of biomolecules can be evaluated with respect to cell response and viability. For example, extracellular vesicles (EVs) substantially influence cultured cell behavior. While all of our products can serve cell response applications, we report here an extensive list of applications in this area for **Cleanascite™ & BindPro™**.


**Cleanascite™** is derived through a proprietary formulation of metallic oxide derivatives. However, unlike other metallic oxides, **Cleanascite™** does not have significant protein binding, making its selectivity profile for lipids unrivaled in the bio-research products industry. As a result, it is ideal to clear lipid-associated matrix effects - including extracellular vesicles, which may influence cell response assays.

#### Removes Lipid Factors

Phospho-Lipids  
>99% Cholesterol &  
Triglycerides  
Lipoproteins  
Extracellular Vesicles  
(Exosomes)



#### Improved Cell Response Performance



- ◆ Proliferation rate factors
- ◆ Cell survival factors
- ◆ Oocyte maturation
- ◆ Toxin Neutralizing Titer/Vaccine

**Cleanascite™** has been validated in accordance with CLIA '88 guidelines. The applications and references for the many diverse investigations using **Cleanascite™** upstream of cell response measurements are next described.

**BindPro™** is a polymeric protein removal suspension reagent, designed to very efficiently remove proteins in an aqueous environment, no solvents are employed.

Chan DW, Yung MM, Chan Y-Sang, Xuan Y, Yang H, Xu D, Zhan J-Biao, Chan KK, Ng T-Bun, Ngan HY, MAP30 protein from *Momordica charantia* is therapeutic and has synergic activity with cisplatin against ovarian cancer in vivo by altering metabolism and inducing ferroptosis, *Pharmacological Research* (2020), doi: <https://doi.org/10.1016/j.phrs.2020.105157>


*Cross-reference classifiers*

Disease: Cancer

Sample Type: Omental conditioned media

Cell Response: Cancer cell lines

Increasing evidence shows that Traditional Chinese Medicine (TCM) has an obvious appeal for cancer treatment, but there is still a lack of scientific investigation of its underlying molecular mechanisms. Bitter melon or bitter gourd (*Momordica charantia*) is an edible fruit that is commonly consumed, and it is used to cure different diseases in various ancient folk medical practices. The investigators report that a bioactive protein, MAP30, isolated from bitter melon seeds exhibited potent anticancer and anti-chemoresistant effects on ovarian cancer cells. To support the analysis of MAP30 altering glycolysis and lipid metabolism in ovarian cancer cells, the authors state "Nile Red fluorescence of lipid-loaded ES2 (human epithelial ovarian cancer cells) in OCM culture for 48 h. ...OCM (Omental conditioned medium) with **Cleanascite™** treatment were used as negative controls.". The authors conclude that natural MAP30 is a non-toxic supplement that may enhance chemotherapeutic outcomes and benefit ovarian cancer patients with peritoneal metastases.



Gomes, A.P., Ilter, D., Low, V. *et al.* Age-induced accumulation of methylmalonic acid promotes tumour progression. *Nature* (2020). <https://doi.org/10.1038/s41586-020-2630-0>

*Cross-reference classifiers*

Disease: Cancer

Sample Type: Human serum

Cell Response: Cancer cell lines

The risk of cancer and associated mortality increases substantially in humans from the age of 65 years onwards. The authors describe how metabolic alterations that occur with age can produce a systemic environment that favors the progression of tumors. Specifically, that methylmalonic acid (MMA), a by-product of propionate metabolism, is upregulated in the serum of older people and functions as a mediator of tumor progression. To support this analysis, the authors state "HS (human serum) samples were manipulated to assess the components ... that might facilitate entrance of MMA into cells. To delipidate the HS, **Cleanascite** Lipid Removal Reagent (Biotech Support Group) was used according to the manufacturer's protocol ..., using a 1:4 volume ratio of

**Cleanascite** reagent to sample". The authors conclude that depletion of lipidic structures from serum taken from an older population, resulted in a reduction in total serum MMA levels and was sufficient to abrogate the pro-aggressive phenotype. The data show that MMA, complexed with lipidic structures, is a circulatory factor that contributes to the pro-aggressive effects of ageing in cancer cells and is sufficient to drive tumour progression and aggressiveness. Thus, MMA is a promising therapeutic target for advanced carcinomas.

Stahl, Elia, et al. "[Phosphatidylcholines from \*Pieris brassicae\* eggs activate an immune response in \*Arabidopsis\*.](#)" eLife 9 (2020): e60293.

*Cross-reference classifiers*


Disease: Plant immune response

Sample Type: *P. brassicae* egg extract

Cell Response: *Arabidopsis* cell lines

Recognition of conserved microbial molecules activates immune responses in plants, a process termed pattern-triggered immunity (PTI). Similarly, insect eggs trigger defenses that impede egg development or attract predators, but information on the nature of egg-associated elicitors is scarce. The authors performed an unbiased bioactivity-guided fractionation of eggs of the butterfly *Pieris brassicae* to determine the nature of egg-associated molecular patterns (EAMPs) that induce immune responses in *Arabidopsis*. Preliminary tests indicated that egg-derived defense eliciting compounds are of lipidic nature. To confirm this, the authors

**A**



stated "we used **Cleanascite** solid-phase aqueous reagent to selectively adsorb lipids from EE (Egg Extract). Application of the lipid-containing phase to ...*Arabidopsis* reporter lines triggered strong and localized GUS staining, similar to EE treatment. In contrast, the supernatant containing proteins and other non-lipidic molecules was not active, indicating that defense gene-inducing molecules were restricted to the lipid phase (Figure 1A)". This helped the research in the identification of phosphatidylcholines (PCs) as egg-associated molecular patterns (EAMPs), and illustrated the

acute ability of plants to detect conserved immunogenic patterns from their enemies, even from seemingly passive structures such as eggs.

Wang, Haiping, et al. "[CD36-mediated metabolic adaptation supports regulatory T cell survival and function in tumors.](#)" *Nature Immunology* (2020): 1-11.

*Cross-reference classifiers*

Disease: Cancer

Sample Type: Conditioned media

Cell Response: Intratumoral Treg cells

Depleting regulatory T cells (Treg cells) to counteract immunosuppressive features of the tumor microenvironment (TME) is an attractive strategy for cancer treatment. However, systemic impairment of their suppressive function limits its therapeutic potential. Elucidating approaches that specifically disrupt intratumoral Treg cells is direly needed for cancer immunotherapy. The use of **Cleanascite**™ helped demonstrate that intratumoral Treg cells increase lipid metabolism and CD36 expression. The article states "cancer cell-conditioned medium ... was treated with **Cleanascite**™ reagent (Biotech Support Group) before Treg cell culture at a volume ratio of 1:5 according to the manufacturer's instructions." The study concludes that CD36 targeting elicited additive antitumor responses with anti-programmed cell death protein 1 therapy. The findings uncover the unexplored metabolic adaptation that orchestrates the survival and functions of intratumoral Treg cells, and the therapeutic potential of targeting this pathway for reprogramming the tumor microenvironment.

Miller-Rhodes, Patrick, and Harris A. Gelbard. [The Cell Culture Environment Regulates the Transcription Factor MafB in BV-2 Microglia.](#) "Matters 7.1 (2021): e202010000001.

*Cross-reference classifiers*

Disease: Central nervous system (CNS)

Sample Type: Conditioned media

Cell Response: BV-2 microglial cells

Microglia experience dramatic molecular and functional changes when transferred from the central nervous system (CNS) to a cell culture environment. This investigation explores lipid dependency as CNS-specific microenvironmental cues, that dictate the gene-regulatory networks specified by master regulator transcription factors such as V-maf musculoaponeurotic fibrosarcoma oncogene homolog B (MafB). To support this analysis, the authors evaluated serum lipid-depletion, stating "Lipids were depleted from FBS using a Cleanascite lipid removal agent (Biotech Support Group, cat no. X2555-10). Cleanascite reagent was thoroughly resuspended before mixing with FBS at a volume ratio of 1:4 (Cleanascite: FBS). The mixture was gently and periodically inverted for 10 min to facilitate lipid binding. The solution was centrifuged at 1,000 x g for 15 min to pellet the removal agent. The resulting supernatant was used for experiments." The report concludes that depletion of lipids, either by serum deprivation or the use of lipid-depleted serum, reduced MafB protein levels in BV-2 microglial cells. In aggregate, the data suggest that serum exposure regulates the transcription factor MafB in BV-2 cells through direct and indirect mechanisms.

Chen, Rain R., et al. "[Targeting of lipid metabolism with a metabolic inhibitor cocktail eradicates peritoneal metastases in ovarian cancer cells.](#)" *Communications Biology* 2 (2019).

*Cross-reference classifiers*

Disease: Cancer

Sample Type: Conditioned media

Cell Response: Ovarian cancer cell lines

Ovarian cancer is an intra-abdominal tumor in which the presence of ascites facilitates metastatic dissemination, and is associated with poor prognosis. However, the significance of metabolic alterations in ovarian cancer cells in the ascites microenvironment remains unclear. In this study, the authors investigated whether reprogramming of lipid metabolism in ovarian cancer cells could modulate cell viability and aggressiveness. The article states: "To determine whether fatty acids in OCM are the primary energy source, fatty acids from OCM was first removed by **Cleanascite™** Lipid Removal Reagent... Then, XTT cell proliferation assays showed that the growth rate of ovarian cancer cells was remarkably reduced in cells cultured in **Cleanascite™**-treated OCM (Fig. 2d). Likewise, co-treatment with **Cleanascite™** and OCM significantly attenuated the increased cell migration and invasion capacities of ES-2 and SKOV3 cells (Fig. 2e, f). These findings suggest that the fatty acid-enriched OCM provides as an energy source for supporting tumor growth and aggressiveness of ovarian cancer cells.". The authors conclude that targeting the lipid metabolism signaling axis impedes ovarian cancer peritoneal metastases.

Turner JD, Langley RS, Johnston KL. [Wolbachia Lipoprotein Stimulates Innate and Adaptive Immunity through Toll-like Receptors 2 and 6 to Induce Disease Manifestations of Filariasis](#) *The Journal of Biological Chemistry*.2009;284:22364-22378


*Cross-reference classifiers*

Disease: Infectious Disease

Sample Type: Soluble *Brugia malayi* female worm extract

Downstream Use Platform: HEK-TLR2 cells

*Wolbachia* endosymbiotic bacteria have been implicated in the inflammatory pathogenesis of filariasis. Inflammation induced by *Brugia malayi* female worm extract (BMFE) is dependent on Toll-like receptors 2 and 6 (TLR2/6) with only a partial requirement for TLR1. Removal of *Wolbachia*, lipids, or proteins eliminates all inflammatory activity. The article states "To determine if TLR2/6 ligands of *Wolbachia* are lipoproteins, we treated the filarial extracts with **Cleanascite™**, which selectively removes lipids and lipoproteins, or with **BindPro™**, a polymeric protein removal suspension reagent (Biotech Support Group). Both treatments completely ablated (to background levels) HEK-TLR2 cell IL-8 reporter gene activity to BMFE (Fig. 1C) thereby showing that the TLR2/6 activity depends on both lipid and protein moieties." The authors conclude that *Wolbachia* lipoproteins drive interferon-dependent CD4<sup>+</sup> T cell polarization and antibody switching.



**The inflammatory stimuli of BMFE are lipoproteins that primarily signal via TLR2/6.** C. triplicate HEK-TLR2 cultures were stimulated with BMFE or control stimuli (doses stated are micrograms/ml) before or following **Cleanascite™** or **BindPro™** treatment. Data plotted are mean IL-8  $\pm$  1S.E.

Lee, Hong-Jai, et al. "[Regulatory effect of humoral milieu on the viral DNA and surface antigen expression of hepatitis B virus \(HBV\) in vitro.](#)" *Molecular & Cellular Toxicology* 15.2 (2019): 123-128.

*Cross-reference classifiers*

Disease: Hepatitis B

Sample Type: Serum

Cell Response: viral DNA and surface antigen expression

The investigations explored if humoral milieu such as serum or culture media, and its constituents, and pH would regulate the viral DNA and surface antigen expression of HBV *in vitro*. Furthermore, lipid removal analysis showed decreased level of HBV DNA and surface antigen expression in human and mouse serum. The article states "To evaluate the lipid exposure status within lipid bilayer, **Cleanascite™** (Biotech Support Group) was added to HBV mixtures in the human serum, mouse serum, or DMEM, and the HBsAg and HBV DNA were evaluated. ... we examined the virus-lipid interaction in non-host milieu, and compared the interaction between in host and non-host milieu. The levels of HBsAg and HBV DNA were significantly decreased with lipid removal by **Cleanascite™** in mouse serum rather than human serum". The authors' concluded that humoral lipid might confer protection to virion against toxicants or hostile interaction with humoral components.

Nguyen, Doan C., et al. "[Extracellular vesicles from bone marrow-derived mesenchymal stromal cells support ex vivo survival of human antibody secreting cells.](#)" *Journal of extracellular vesicles* 7.1 (2018): 1463778.

*Cross-reference classifiers*

Disease: Pan/Methods

Sample Type: Bone marrow-derived mesenchymal stromal cell secretome

Cell Response: Human antibody secreting cells

Bone marrow-derived mesenchymal stromal cells (MSC) have been shown to support human antibody secreting cells (ASC) survival *ex vivo*. Extracellular vesicles from bone marrow-derived mesenchymal stromal have novel mechanisms of cell-cell communication over short and long distances, but whether the crosstalk between these cell interactions can occur via extracellular vesicles is not known. Thus, the study investigated the role of extracellular vesicles (EVs) in antibody secreting cell survival and IgG secretion. The article states "To understand whether disrupting the lipid plasma membrane that upsets the integrity of the EVs would compromise the survival activity of either the non-irradiated or irradiated secretomes, ... We then cultured ASC with conventional media (vehicle), secretome from irradiated MSC, or secretome from irradiated MSC that had been pretreated with the lipid-disrupting agent **Cleanascite™** [28], which is known not to alter protein functionality [29]. **Cleanascite™**-treatment of the secretome dramatically reduced ASC functional survival, ... Similar reductions were also noted with the secretome of non-irradiated MSC when treated with **Cleanascite™** ... These results demonstrate that lipid-membrane bodies, such as EVs, could mediate important ASC survival factors within the MSC secretome."

Sprenkle, Neil T., et al. "[Endoplasmic reticulum stress is transmissible in vitro between cells of the central nervous system.](#)" *Journal of Neurochemistry*.

*Cross-reference classifiers*

Disease: Neurodegenerative

Sample Type: Conditioned media

Cell Response: Stressed astrocytes and neurons



Improper protein folding and trafficking are common pathological events in neurodegenerative diseases that result in the toxic accumulation of misfolded proteins within the lumen of the endoplasmic reticulum (ER). The cell-extrinsic role of sustained unfolded protein response activation under physiological and pathological states in the central nervous system (CNS) remains to be elucidated. The authors studied the characteristics of a mediator secreted by ER stressed astrocytes and neurons. To determine if the mediator was a lipid associated factor, the article states "...100 µl of **Cleanascite™** slurry was added to 1 ml of conditioned medium and incubated at RT with end-over-end mixing for 1 h followed by centrifugation." The authors provided evidence that depletion of lipids from astrocyte conditioned media using **Cleanascite™** abrogated transmission of ER stress. Such evidence helped the authors conclude that ER stressed astrocytes and neurons secrete a molecule(s) with lipid characteristics which regulates both inflammatory and ER stress responses in other astrocytes, neurons, and microglia *in vitro*. These findings provide insight into the cell-nonautonomous influence of ER stress on cells of the central nervous system.

Barrera N, dos Santos Neto PC, Cuadro F, Bosolasco D, Mulet AP, Crispo M, et al. (2018) Impact of delipidated estrous sheep serum supplementation on in vitro maturation, cryotolerance and endoplasmic reticulum stress gene expression of sheep oocytes. PLoS ONE 13(6): e0198742. <https://doi.org/10.1371/journal.pone.0198742>

*Cross-reference classifiers*

Disease: Cryosurvival

Sample Type: Estrous sheep serum

Cell Response: Oocyte maturation

High lipid content of oocytes and embryos in domestic animals is one of the well-known factors associated with poor cryosurvival. In this articles, the authors wanted to determine whether the use of delipidated estrous sheep serum during in vitro maturation (IVM) of ovine oocytes reduces the cytoplasmic lipid droplets content and improves embryo development and cryotolerance after vitrification. The article states "Lipid removal from serum was performed by using **Cleanascite™** (Biotech Support Group, NJ, USA) according to the instructions provided by the manufacturer. Unlike other approaches, the protocol described herein for delipidation of estrous sheep serum was effective in decreasing levels of Triglycerides, total Cholesterol, and NEFAs. To our knowledge this is the first study to use the **Cleanascite™** method to generate estrous sheep serum yielding significantly reduced lipid levels. Subsequent use of the partially delipidated serum as supplemented in IVM media resulted in effective reduction of oocyte lipid content. The advantage of this method over other traditional methods (i.e. chloroform) includes increased feasibility and reduced toxicity and biosafety concerns". Their results demonstrate that although supplementation of IVM medium with delipidated estrous sheep serum reduces the presence of cytoplasmic lipid droplets in oocytes after maturation, oocyte cryotolerance is not improved.

Dean, E. Danielle, et al. "[Interrupted glucagon signaling reveals hepatic  \$\alpha\$  cell axis and role for L-glutamine in  \$\alpha\$  cell proliferation.](#)" Cell metabolism 25.6 (2017): 1362-1373.

*Cross-reference classifiers*

Disease: Diabetes

Sample Type: Mouse serum

Cell Response:  $\alpha$  cell proliferation

Decreasing glucagon action lowers blood glucose and may be useful therapeutically for diabetes. However, interrupted glucagon signaling leads to  $\alpha$  cell proliferation. In this article, the authors wanted to determine which factors affected  $\alpha$  cell proliferation. The article states "For lipid removal, whole mouse serum was treated with **Cleanascite™** reagent (Biotech Support Group, Monmouth Junction, NJ) prior to islet culture at a 1:1 ratio according to the vendor's protocol. Lipid removal was validated by HPLC to remove 99% of all phospholipids, cholesterols, and triglycerides....". In testing whether lipids could stimulate  $\alpha$  cell proliferation, it was found that serum activity was retained after the removal of >99% of triglycerides, cholesterols, and phospholipids. The authors conclude that amino acids, especially L-glutamine, regulate  $\alpha$  cell proliferation and mass via mTOR-dependent nutrient sensing.

Lovászi, M., et al. "[Sebum lipids influence macrophage polarization and activation.](#)" *British Journal of Dermatology* (2017). doi: 10.1111/bjd.15754.

*Cross-reference classifiers*

Disease: Cell Biology

Sample Type: Sebocytes

Downstream Use Platform: Immortalized human SZ95 sebocytes

The article's authors report on sebum lipids contributing to the differentiation, polarization and function of macrophages. In order to determine the role of specific lipids, lipid removal was investigated from supernatants of the immortalized human SZ95 sebocytes, as stated, "For lipid depletion of the supernatants **Cleanascite™** lipid clarification reagent (Biotech Support Group, Monmouth Junction, NJ, USA) was used according to the manufacturer's instructions. Lipids; squalene, linoleic acid, oleic acid, palmitic acid and stearic acid (Sigma-Aldrich); were replaced individually subsequent to lipid depletion in a concentration of 150 µM.". The authors concluded a role for sebaceous glands in modulating immune responses via their secreted lipids that are of possible pathologic and therapeutic relevance.

Chan, DW, Mak, SL, Ngan, HYS. The significance of lipid metabolism in peritoneal metastases of ovarian cancer. The 2016 Cold Spring Harbour Asia Conference on Cancer and Metabolism, Suzhou, China, 19-23 September 2016. <http://hub.hku.hk/handle/10722/235385>

*Cross-reference classifiers*

Disease: Cancer

Sample Type: Omental explant culture

Downstream Use Platform: Ovarian oncogenic capacities

The authors report that the high lipid content in ascetic fluid provides a huge energy source for ovarian cancer cells in peritoneal dissemination and intraperitoneal tumor colonization. In this study, ovarian cancer cells co-cultured with an omental explant culture system (OCM) or ascetic fluid from ovarian cancer patients exhibited an increase in *in vitro* cell growth, cell migration/invasion through activation of TAK1/NF-kappaB signaling cascade. The abstract states "In contrast, the oncogenic capacities of ovarian cancer cells were impaired when cultured in OCM treated with **Cleanascite™** Lipid Removal Reagent, suggesting that the bioactive lipids in OCM are required for enhanced oncogenic capacities".

Kamtchoua, Thierry, Monica Bologna, Robert Hopfer, David Neveu, Branda Hu, Xiaohua Sheng, Nicolas Corde, Catherine Pouzet, Gloria Zimmerman, and Sanjay Gurunathan. [Safety and immunogenicity of the pneumococcal pneumolysin derivative PlyD1 in a single-antigen protein vaccine candidate in adults](#). Vaccine (2012).

*Cross-reference classifiers*

Disease: Infectious Disease Vaccine

Sample Type: Serum

Downstream Use Platform: toxin-neutralizing antibody titer

Authors Thierry Kamtchoua et al published an article in the journal Vaccine titled, 'Safety and immunogenicity of the pneumococcal pneumolysin derivative PlyD1 in a single-antigen protein vaccine candidate in adults' describing the immunogenicity of pneumococcal single antigen protein vaccine in a phase 1, randomized, placebo controlled dose escalating study. Authors cite **Cleanascite™** from Biotech Support Group for removal of cholesterol from serum. A toxin neutralizing assay with antibodies in sera was developed to neutralize cytotoxicity caused by Ply in Vero cells. An incubated challenge dose of pneumolysin toxin containing serum diluted with or without **Cleanascite™** was developed. The neutralizing titer inhibited the toxin's effect on Vero cells. According to the paper, "Briefly, the toxin-neutralizing antibody titer was determined by incubating a challenge dose of pneumolysin toxin with serial 2-fold dilutions of serum treated with or without **Cleanascite™** (Biotech Support Group) to remove cholesterol, an inhibitor of Ply".

Antunes RF; Brandao C; Maia M; Arosa FA. [Red blood cells release factors with growth and survival bioactivities for normal and leukemic T cells](#). Immunology and Cell Biology.2011;89(1):111-21

*Cross-reference classifiers*

Disease: Cancer

Sample Type: Red Blood Cell media

Downstream Use Platform: T Cell proliferation

Red blood cells (RBCs) have been implicated since the early 1970s in the modulation of T cell responses both *in vitro* and *in vivo*. As they can also regulate biological processes of neighboring cells, the authors aim was to show that human red blood cell conditioned media contains bioactive factors that favor proliferation of normal activated T cells and leukemic Jurkat T cells. To define whether factors that favor proliferation were lipid associated, the RBC media was depleted of extracellular vesicles by ultracentrifugation. Then, the article states "For lipid depletion, **Cleanascite™** was added to the RBC-sup in a ratio 1:4 and the mixture incubated first in a rotator at room temperature for 10 min, followed by a further incubation at 4°C for 30 min, following manufacturer's

instructions. Then, the mixture was centrifuged to remove the resin and the RBC-sup collected and concentrated as indicated above before the *in vitro* bioactivity assays". The authors conclude that red blood cells release protein factors (not lipid factors) with the capacity to sustain T-cell growth and survival. Such protein factors may have an unforeseen role in sustaining malignant cell growth and survival *in vivo*.

Cheng AM, Moore EE, Masuno T et al [Normal Mesenteric Lymph Blunts the Pulmonary Inflammatory Response to Endotoxin](#). Journal of Surgical Research.2006;136(S2):166-171

*Cross-reference classifiers*

Disease: Hemorrhagic shock

Sample Type: Mesenteric lymph

Downstream Use Platform: Primary human pulmonary endothelial cells

LPS induced ICAM-1 expression decreases by lipoproteins in normal mesenteric lymph(NML) which contain anti-inflammatory factors. **Cleanascite™** was used for delipidation and removal of lipoproteins from primary human pulmonary endothelial cells (HMVECs) incubated with normal mesenteric lymph NML or post-shock mesenteric lymph PSML. ICAM expression was measured after LPS stimulation by flow cytometry. ICAM-1 surface expression was measured by flow cytometry. **Cleanascite™** extracted lipoproteins from NML before incubation and LPS-induced ICAM-1 expression was determined. Researchers concluded that decreased lipoprotein expression after hemorrhagic shock HS increases post-shock mesenteric lymph PSML toxicity from the ischemic gut.

Cho N, Chueh PJ, Kim C et al [Monoclonal antibody to a cancer-specific and drug-responsive hydroquinone \(NADH\) oxidase from the sera of cancer patients](#). Cancer Immunology, Immunotherapy. 2002;51(3):121-9

*Cross-reference classifiers*

Disease: Cancer

Sample Type: Serum

Downstream Use Platform: Cancer cell cultures

Scientists prepared monoclonal antibodies to a 34-kDa circulating form of a drug-responsive hydroquinone (NADH) oxidase with a protein disulfide–thiol interchange activity specific to the surface of cancer cells and the sera of cancer patients. **Cleanascite™** was used for delipidation of sera. Epitopes (antibody (mAb) 12.1 and postimmune antisera ) inhibited the drug-responsive oxidation of NADH with the sera of cancer patients. Authors concluded both mouse ascites containing mAb 12.1 and postimmune sera (but not preimmune sera) slowed the growth of human cancer cell lines in culture, but did not affect the growth of non-cancerous cell lines.

## Patents (Third Party)

United States Patent Application 20170348398 entitled: "[COMPOSITIONS AND METHODS FOR DECREASING BLOOD GLUCAGON LEVELS](#)"

*Cross-reference classifiers*

Disease: Diabetes

Sample Type: Serum

Cell Response: Pancreatic  $\alpha$  cell proliferation

The patent discloses compositions and methods for decreasing blood glucagon levels. As disclosed, L-glutamine is a selective stimulator of  $\alpha$ -cell proliferation generated when glucagon signaling is interrupted. A method for treating a subject with hyperglucagonemia, e.g., a subject with diabetes, that involves administering to the subject a composition comprising an L-glutamine inhibitor in an amount effective to decrease blood glucagon levels, is disclosed. In an example, pancreatic islets were isolated from male 8-14 week old C57B16/J mice (Jackson Laboratory, ME) and cultured in various media conditions for 3 days. The patent states " For lipid removal, serum was treated with **Cleanascite™** reagent (Biotech Support Group, Monmouth Junction, N.J.) prior to islet culture at a 1:1 ratio according to the vendor's protocol. ". The example supports that increased amino acids, but not lipids and other soluble factors, selectively increased rapamycin-sensitive  $\alpha$ -cell proliferation.



## Cell Response Products

- Extensively cited
- Replaces hazardous hydrocarbons
- Diverse samples

### Cleanascite™

#### *Lipid Adsorption & Clarification*

- Effectively replaces chlorinated/fluorinated hydrocarbons (eg. freon)
- Workflows for antibodies, proteins, nucleic acids, proteoglycans, and most serum analytes
- A high binding capacity for lipids with minimal cross-reactivity with proteins and nucleic acids
- Ideal for clarifying ascites, serum, cell & tissue culture, bile and organ homogenates
- Compatible with cell and tissue culture
- Exquisite selectivity profile including extracellular vesicle and exosome clearance
- Enrichment of delipidated tissue samples

### BindPro™

#### *Aqueous Protein Crash & Enrichment of Metabolites/Analytes*

- Serum and plasma protein removal, >95%
- Aqueous buffer system, compatible with cell and tissue culture
- Aqueous Protein Crash, linearly scalable, unlike chemical precipitation or membrane filtration.
- Applicable for drug binding/screening, target analytes and metabolomics
- Protein removal is species agnostic; sera tested includes human, mouse, sheep, bovine, goat, rat

**Cleanascite™** and **BindPro™** are supplied as suspension reagents. Simply add, mix and centrifuge in a 10 minute protocol.

5 ml of each reagent can be purchased as a trial kit.

Product	Size	Item No.	Price
<b>Cell Response Trial Kit</b>	<b>BindPro™</b> 5 ml	CRT100-5	\$325
	<b>Cleanascite™</b> 5 ml		



[www.biotechsupportgroup.com](http://www.biotechsupportgroup.com)